

# Review of the genus *Coccus* Linnaeus from Korea, with description of a new species (Hemiptera, Coccoomorpha, Coccidae)

Jinyeong Choi<sup>1,2</sup>, Seunghwan Lee<sup>1,2</sup>

**1** *Insect Biosystematics Laboratory, Department of Agricultural Biotechnology, Seoul National University, Seoul 151-921, Republic of Korea* **2** *Research Institute of Agriculture and Life Sciences, Seoul National University, Seoul 151-921, Republic of Korea*

Corresponding author: *Seunghwan Lee* ([seung@snu.ac.kr](mailto:seung@snu.ac.kr))

---

Academic editor: *R. Blackman* | Received 5 December 2017 | Accepted 20 December 2011 | Published 5 February 2018

---

<http://zoobank.org/2FD0E47C-9C5C-4022-ABF7-8482872414A0>

---

**Citation:** Choi J, Lee S (2018) Review of the genus *Coccus* Linnaeus from Korea, with description of a new species (Hemiptera, Coccoomorpha, Coccidae). ZooKeys 734: 121–135. <https://doi.org/10.3897/zookeys.734.22774>

---

## Abstract

The genus *Coccus* from Korea is reviewed, including a new species, *C. fvicola* **sp. n.**, and a first record of *C. malloti* (Takahashi, 1956). The new species is characterized by a distinctive pattern of ventral tubular ducts on medial area of head and thorax, and submarginal area of abdomen. The adult female of *Coccus fvicola* **sp. n.** is described and illustrated, and a morphological comparison is given with congeners. *Coccus malloti* is redescribed and illustrated based on the adult female specimens from Korea. A key to the four species of *Coccus* known from Korea is provided with diagnoses and photographs.

## Keywords

Coccinae, Coccini, soft scale insect, taxonomy

## Introduction

The genus *Coccus* Linnaeus, 1758, which is a species-rich group in the family Coccidae, comprises approximately 111 species worldwide (Hodgson 1994; García-Morales et al. 2016). This group is defined morphologically by the distribution of ventral tubular ducts, the shape of dorsal and marginal setae, and the presence of a tibio-tarsal articulatory sclerosis on each leg (Hodgson 1994); however, the molecular phylogeny using

some of the taxa included in Coccidae revealed that it is not a monophyletic group and needs taxonomic revision (Lin et al. 2013). Among the genus *Coccus*, *C. hesperidum* Linnaeus, *C. viridis* Green, and *C. celatus* De Lotto are known as economic pests of ornamental and agricultural products (Kapranas et al. 2007; Waller et al. 2007). Two species of the genus, *C. hesperidum* and *C. pseudomagnoliarum* (Kuwana) have been previously recorded from Korea. Here, a new species is described, *Coccus ficicola* sp. n. and its morphology is compared with congeners. A hitherto unrecorded species is also redescribed, *C. malloti* (Takahashi) and recorded for the first time from Korea.

## Materials and methods

The specimens were mounted on microscope slides using the method of Hodgson and Henderson (2000) and Danzig and Gavrilov-Zimin (2014). The micrographs of slide-mounted materials were taken and measured using analysis software (Active Measure ver. 3.0.3, Mitani Co. Ltd, Japan). The terminology follows Hodgson (1994) and Hodgson and Henderson (2000), except that the term “pregenital disc-pores” is replaced with “multilocular pores” suggested by Kondo and Hardy (2008). The type specimens are deposited in the Insect Biosystematics Laboratory, Research Institute for Agriculture and Life Science, Seoul National University, Korea (SNU).

## Taxonomy

### Genus *Coccus* Linnaeus, 1758: 455

**Type species.** *Coccus hesperidum* Linnaeus, 1758, designated by Opinion 1303 (1985).

**Diagnosis.** Dorsal setae pointed or blunt; dorsal tubular ducts and dorsal tubercles present or absent; marginal setae with pointed or frayed apices; ventral tubular ducts present or absent, if present, mainly distributed on medial area of thorax or submarginal area; a tibio-tarsal articular sclerite present or absent on each leg. For further diagnostic characteristics, see Hodgson (1994) and Wang and Feng (2012).

### Key to species of genus *Coccus* in Korea

- 1 Dorsal tubercles absent; legs without tibio-tarsal articular sclerites; ventral tubular ducts present on abdomen only.... *C. pseudomagnoliarum* (Kuwana)
- Dorsal tubercles present; legs with tibio-tarsal articular sclerites; ventral tubular ducts present on thorax and abdomen ..... **2**
- 2 Ventral tubular ducts of three types (Type I: each with a broad inner ductule; Type II: each with a narrow inner ductule; Type III: each with a filamentous inner ductule) present..... *C. malloti* (Takahashi)
- Ventral tubular ducts of type I, each with a narrow inner ductule ..... **3**

- 3 Antenna 7-segmented; ventral tubular ducts scarce: a small group of 0–3 ducts present between mouthparts and each procoxa; a thin transverse band containing one or two ducts vertically present between mesocoxae; absent on inner submarginal area of abdomen ..... *C. hesperidum* (Linnaeus)
- Antenna 8-segmented; ventral tubular ducts abundant: a large group of 16–20 ducts present between mouthparts and each procoxa; a broad transverse band containing 4–7 ducts present vertically between mesocoxae; present on inner submarginal area of abdomen ..... *C. ficicola* sp. n.

***Coccus ficicola* sp. n.**

<http://zoobank.org/1C880C8E-9D53-4418-A87D-EFBBB2227497>

Figs 1A–D, 2A–Q


**Material examined. Holotype:** adult female: Korea, Gangnam-gu, Yeoksam-dong, 18.iv.2015, coll. J.Y. Choi, on *Ficus benghalensis* L. (Moraceae). **Paratypes:** same data as holotype, 9♀♀.

**Diagnosis.** Adult females in life (Fig. 1A–D) with a reticulated pattern of brown stripes and a longitudinal ridge medially on dorsum; dermal areolations present but small; dorsal tubercles present; dorsal tubular ducts sparse on submarginal area; dorsal setae with bluntly rounded apices; marginal setae usually with simple pointed apices; multilocular disc-pores usually with ten loculi; ventral tubular ducts with a narrow inner ductule, frequent on posterior region of the head, medial area of thorax, and inner submarginal area of abdomen; antennae each with eight segments; legs each with a tibio-tarsal sclerosis on the articulation.

**Description. Adult female. Living appearance** (Fig. 1A–D). Body oval, flattened, or moderately convex. Young adult females yellowish to brownish, with a reticulated pattern of brown stripes except for a longitudinal ridge on mid dorsum. Older adult females becoming more convex and darker. Eggs not seen.

**Slide-mounted material** (Fig. 2A–Q). Body oval, 2.6–3.5 mm long, 2.0–3.6 mm wide, with distinct stigmatic cleft; anal clefts approximately 1/6 of body length.

*Dorsum.* Derm membranous. Dermal areolations oval and small, each with a microduct. Dorsal tubercles normally convex, present on submarginal area, 4–6 in total on each side: two between apex of head and anterior stigmatic cleft, one or two between anterior and posterior stigmatic clefts, and one or two between posterior stigmatic cleft and anal cleft. Dorsal setae cylindrical, short, stout, blunt apically, each 6–9 µm long, moderately distributed on dorsum. Dorsal tubular ducts each with a developed outer ductule and a slender inner ductule with a developed terminal gland, sparsely present on submargin. Dorsal microducts evenly scattered over entire dorsum. Preopercular pores round and small, rather inconspicuous, 6–7 µm wide, set in a small group of approximately 6–15 in front of anal plates. Anal plates quadrate, 190–223 µm long, 160–203 µm wide, usually posterolateral margin slightly longer than anterolateral margin; anterolateral margin 119–144 µm long, posterolateral margin 130–154 µm




**Figure 1.** *Coccus ficicola* Choi & Lee sp. n. **A** instar female **B** immature adult female **C** mature adult female **D** mature (upper) and immature (under) adult females.

long; each plate with four apical setae. Ano-genital fold with two pairs of anterior and three pairs of lateral margin setae. Anal ring with six long setae.

*Margin.* Marginal setae spinose, slender, slightly bent, each 16–32  $\mu\text{m}$  long, mostly with simple pointed apices, but sometimes with bifid tips; with 52–59 present between anterior stigmatic clefts; 12–17 laterally present between anterior and posterior stigmatic clefts, 45–53 present between posterior stigmatic cleft and anal clefts. Stigmatic clefts deep, each with three stigmatic spines, median spine 2–3 times as long as lateral spine: medians 48–77  $\mu\text{m}$  long, laterals 14–30  $\mu\text{m}$  long. Eyespots located near margin.

*Venter.* Derm membranous. Multilocular disc-pores 7–8  $\mu\text{m}$  wide, each with 10–12 loculi, mostly with ten loculi, abundant around vulvar area, but less frequent on anterior segments of abdomen. Spiracular pores 4–5  $\mu\text{m}$  wide, each with five loculi, in a narrow band 1–2 pores wide between each spiracle and stigmatic cleft. Ventral tubular ducts of a single type, each 21–28  $\mu\text{m}$  long, with a developed outer ductule and a narrow inner ductule with a flower-shaped terminal gland, approximately 16–20 ducts densely present between mouthparts and procoxa on each side; a broad transverse band containing around 4–7 ducts vertically present between mesocoxae; abundant between each meso- and metacoxa, extending around spiracles; and also sparsely scattered on



**Figure 2.** *Coccus ficicola* sp. n., adult female. **A** dorsal microduct **B** dorsal tubercle **C** dermal areolations **D** dorsal seta **E** stigmatic spines **F** marginal setae **G** dorsal tubular duct **H** anal plates **I** preopercular pore **J** multilocular disc-pores **K** ventral tubular duct on abdomen **L** leg **M** ventral seta **N** ventral tubular duct on head and thorax. **O** spiracular pore **P** ventral microduct **Q** antenna. Scale bars: 200  $\mu\text{m}$  (**L**, **Q**); 100  $\mu\text{m}$  (**H**); 50  $\mu\text{m}$  (**C**, **E**); others = 10  $\mu\text{m}$ .



inner submarginal area between anal plates and each metacoxa. Ventral microducts present on entire venter, especially frequent on submargin. Ventral setae with three pairs of long pregenital setae; two pairs of long setae between antennae; other setae sharply spinose, each 7–14  $\mu\text{m}$  long, moderately distributed over entire venter. Legs well developed, each with a tibio-tarsal articulation and an articular sclerite; total length of each metathoracic leg 560–638  $\mu\text{m}$  long: each coxa 138–163  $\mu\text{m}$  long, trochanter+femur 181–213  $\mu\text{m}$  long, tibia+tarsus 218–237  $\mu\text{m}$  long, claw 17–26  $\mu\text{m}$  long. Tarsal digitules thinner and longer than claw digitules. Spiracles normal, mostly posterior peritreme broader than anterior: anterior peritremes each 38–49  $\mu\text{m}$  wide, posterior peritremes each 45–57  $\mu\text{m}$  wide. Antenna 8-segmented, each 279–339  $\mu\text{m}$  long. Clypeolabral shield 127–138  $\mu\text{m}$  wide.

**Etymology.** Named after its host plant, *Ficus benghalensis* L.

**Host plant.** Moraceae: *Ficus benghalensis* L.

**Comments.** *Coccus ficicola* sp. n. is probably a non-endemic species because it occurs on an imported ornamental plant, *Ficus benghalensis*, which is widely cultivated in tropical areas (Starr et al. 2003). In order to know the exact origin of the new species, further investigations are needed.

**Morphological comparison of adult females of *Coccus ficicola* sp. n. and its related taxa.** Based on taxonomic articles, such as Gill et al. (1977), Ben-Dov (1981), Avasthi and Shafee (1991), and Lin et al. (2017), we selected ten species morphologically similar to *C. ficicola* sp. n.: *C. capparidis* (Green, 1904), *C. discrepans* (Green, 1904), *C. elatensis* (Ben-Dov, 1981), *C. formicarii* (Green, 1896), *C. gymnospori* (Green, 1908), *C. hesperidum* (Linnaeus, 1758), *C. latioferulatum* (Green, 1922), *C. moestus* (De Lotto, 1959), *C. praetermissus* Lin & Tanaka, 2017, and *C. sulawesicus* Gavrilov, 2013. The morphological characters of adult females of *Coccus ficicola* and the ten species are summarized in Table 1.

In the morphological comparison, *Coccus ficicola* shows a new combination of morphological characters; in particular, the distributional pattern of ventral tubular ducts of the species reveals uniqueness among the nine morphological characters. *Coccus ficicola* is most closely related to *C. gymnospori* (Green), in having (i) dorsal tubercles, (ii) dorsal tubular ducts on submarginal area, (iii) dorsal setae with bluntly rounded apices, (iv) preopercular pores, (v) marginal setae with pointed or frayed apices, (vi) antenna with eight segments, (vii) three pairs of pregenital setae, and (viii) tibio-tarsal sclerite. However, *C. ficicola* differs from *C. gymnospori* in having the following combination of character states (character states of *C. gymnospori* in parenthesis): (i) ventral tubular ducts abundant, 16 to 20 ducts present between mouthparts and each procoxa (few, only 3 or 4 ducts); a broad transverse band containing 4–7 ducts vertically between metacoxae (thin, containing one or two ducts); and present on inner submarginal area of abdomen (entirely absent), and (ii) multilocular disc-pores extending further anteriorly (restricted to preceding two abdominal segments) (Ben-Dov 1981; Avasthi and Shafee 1989).

Although the African species, *C. africanus* (Newstead) and *C. alpinus* De Lotto, are not included in the list of related taxa for morphological comparison, *C. ficicola* is

**Table 1.** Comparison of morphological characters of adult females of *Coccus ficticola* sp. n. and its related taxa.

Species	Dorsal tubercles	Dorsal tubular ducts	Dorsal setae	Preopercular pores	Marginal setae	Antenna	Pregenital setae	Tibio-tarsal sclerotosis	Ventral tubular ducts	Reference
<i>C. ficticola</i> sp. n.	Present	Present	Bluntly rounded	Present	Pointed or frayed	Eight segments	Three pairs	Present	Present on medial area of head, pro- and mesothorax; submarginal area of abdomen	This study
<i>C. capparisidis</i>	Present	Absent	Bluntly rounded	Present	Pointed or frayed	Six or seven segments	One or two pairs	Absent	Present on submarginal area of abdomen	Williams and Watson 1990
<i>C. discrepans</i>	Present	Absent	Sharply pointed or bluntly rounded	absent	Pointed or frayed	Seven segments	Three pairs	Present	Present on medial area of mesothorax	Avasthi and Shafee 1991; Tao et al. 1983
<i>C. elatensis</i>	Present	Present	Bluntly rounded	Present	Pointed or frayed	Eight segments	Four pairs	Present	Present on medial area of mesothorax	Ben-Dov 1981
<i>C. fornicarii</i>	Absent	Absent	Sharply pointed (setose)	Present	Pointed (setose)	Seven or eight segments	Three pairs	Absent	Present on medial area of head, pro- and mesothorax	Hodgson 1994
<i>C. gymnospori</i>	Present	Present	Bluntly rounded	Present	Pointed or frayed	Eight segments	Three pairs	Present	Present on medial area of head and mesothorax	Ben-Dov 1981
<i>C. hesperidum</i>	Present	Present or absent	Sharply pointed	Present	Pointed or frayed	Seven segments	Three pairs	Present	Present on medial area of pro- and mesothorax; laterad to genital opening	Hodgson 1994; Lin et al. 2017
<i>C. latiooperculatum</i>	Absent	Absent	Bluntly rounded	Present	Frayed	Seven segments	Two pairs	Present	Present on medial area of pro- and mesothorax	Avasthi and Shafee 1991
<i>C. moestis</i>	Present	Present	Bluntly rounded	Present	Frayed	Seven or eight segments	Three pairs	Present	Present on medial area of mesothorax	Gill et al. 1977
<i>C. praetermissus</i>	Present	Present	Bluntly rounded	Present	Pointed or frayed	Seven segments	Three pairs	Present	Present on medial area of mesothorax	Lin et al. 2017
<i>C. sulawesicus</i>	Absent	Absent	Sharply pointed	Present	Pointed or frayed	Seven or eight segments	Three pairs	Present	Present on medial area of meso- and metathorax	Gavrilov-Zimin 2013

similar to both species in having abundant ventral tubular ducts. However, *C. ficicola* does not have continuous ventral tubular ducts between the metacoxae, whereas both African species have this character state (De Lotto 1957; De Lotto 1960; Granara de Willink et al. 2010).

### ***Coccus hesperidum* (Linnaeus, 1758)**

Fig. 3A–F

*Coccus hesperidum* Linnaeus, 1758: 455.

**Material examined.** Adult female: Daehak-dong, Gwanak-gu, Seoul, 09.iv.2014, coll. J.Y. Choi, on Orchidaceae sp., 5♀♀; Sinhyo-dong, Seogwipo-si, Jeju-do, 14.ix.2014, coll. J.Y. Choi, on *Asplenium antiquum* Makino (Aspleniaceae), 5♀♀; Sinbuk-eup, Chuncheon-si, Gangwon-do, 31.v.2015, coll. J.Y. Choi, on *Heteropanax fragrans* (Roxb.) (Araliaceae), 5♀♀; Geumam-dong, Deokjin-gu, Jeonju-si, Jeollabuk-do, 06.vi.2015, on same host, 5♀♀; Songhyeon-dong, Andong-si, Gyeongsangbuk-do, 07.vi.2015, on same host, 5♀♀; Guseo-dong, Geumjeong-gu, Busan, 07.vi.2015, on *Ficus benghalensis* L., 5♀♀.

**Diagnosis.** Adult females in life (Fig. 3A–F) highly variable in body color and pigment pattern, but usually dorsum pale yellowish to brownish, with black or brown spots; dermal areolations present; dorsal tubercles present; dorsal tubular ducts present or absent; dorsal setae with sharply pointed apices; marginal setae usually with pointed, bifid or fimbriate apices; multilocular disc-pores usually with ten loculi; ventral tubular ducts with a narrow inner ductule, few present around meso- and procoxa, and anal plates; antennae each 7-segmented; legs each with a tibio-tarsal articulatory sclerosis.

**Host plant.** Recorded from 346 genera in 121 families (García-Morales et al. 2016). For Korean records, see Paik (1978).

**Distribution.** Known from all zoogeographical regions (García-Morales et al. 2016).

### ***Coccus malloti* (Takahashi, 1956)**


Figs 4A–B, 5A–Q

*Pulvinaria malloti* Takahashi, 1956: 25.


**Material examined.** Adult female: Jeollanam-do, Gwangyang-si, Ongnyong-myeon, Chusan-ri, 28.v.2015, coll. J.Y. Choi, on *Ilex cornuta* Lindl. (Aquifoliaceae), 9♀♀; Jeju-do, Seogwipo-si, Andeok-myeon, Gamsan-ri, 27.iv.2016, coll. J.Y. Choi, on *Aphananthe aspera* (Thunb.) (Cannabaceae), 1♀.

**Diagnosis.** Adult females in life (Fig. 4A–B) with a reticulated pattern of black stripes and a longitudinal band medially on dorsum; dermal areolations present but small; dorsal tubercles present; dorsal tubular ducts absent; dorsal setae sharply spinose; marginal setae mostly with simple pointed apices; multilocular disc-pores usually with





**Figure 3.** *Coccus hesperidum* (Linnaeus, 1758). **A** population on *Asplenium antiquum* **B** adult female on *Ficus benghalensis* **C** adult female on Orchidaceae sp. **D, E, F** populations on *Heteropanax fragrans*.



**Figure 4.** *Coccus malloti* (Takahashi, 1956). **A** adult female on *Ilex cornuta* **B** adult female on *Aphananthe aspera*.

ten loculi; ventral tubular ducts of three types: Type I with a broad inner ductule, densely present on posterior medial area of head; frequent on anterior medial area of prothorax, extending to inner submarginal area of thorax; and also sparsely scattered on inner submarginal area of abdomen; Type II with a narrow inner ductule and a large flower-shaped terminal gland, rarely present on inner submarginal area and posterior medial area of abdomen; Type III with a long filamentous inner ductule and a quite small terminal gland, moderately present on submarginal area between anal clefts and each posterior spiracular furrow; all types of ventral tubular ducts absent on medial area of meso-, metathorax and anterior abdomen, and submarginal area of head; antennae each eight segments; legs each with a tibio-tarsal sclerosis on the articulation.


**Redescription. Adult female. Living appearance** (Figs 4A–B). Body elongate oval, flattened, or slightly convex. Young adult females yellowish to dark brownish, with a reticulated pattern of brown or black stripes, getting darker at maturity. Eggs reddish in color, stored beneath venter.

**Slide-mounted material** (Fig. 5A–Q). Body elongate oval, 3.6–4.8 mm long, 2.2–3.3 mm wide, with shallow to deep stigmatic cleft; anal clefts approximately 1/6 of body length.

**Dorsum.** Derm membranous. Dermal areolations oval and small, each with a microduct. Dorsal tubercles normally convex, present on submarginal area, 1–5 in total on each side: one or two between apex of head and anterior stigmatic cleft, zero to two between anterior and posterior stigmatic clefts, and zero or one between posterior stigmatic cleft and anal cleft. Dorsal setae sharply spinose, short, stout, each 6–9  $\mu\text{m}$  long, moderately distributed on dorsum. Dorsal tubular ducts absent. Dorsal microducts evenly scattered over entire dorsum. Preopercular pores round and small, 3–5  $\mu\text{m}$  wide, set in a small group of approximately 8 in front of anal plates. Anal plates quadrate, 217–249  $\mu\text{m}$  long, 169–198  $\mu\text{m}$  wide, each with slightly concaved posterolateral margin, usually posterolateral margin quite longer than anterolateral margin; anterolateral margin 123–143  $\mu\text{m}$  long, posterolateral margin 150–169  $\mu\text{m}$  long; each plate with four apical setae. Ano-genital fold with two pairs of anterior and three pairs of lateral margin setae. Anal ring with six long setae.

**Margin.** Marginal setae spinose, straight or slightly bent, each 14–22  $\mu\text{m}$  long, mostly with simple pointed apices; with 30–43 present between anterior stigmatic clefts; 13–17 laterally present between anterior and posterior stigmatic clefts, 22–38 present between posterior stigmatic cleft and anal clefts. Stigmatic clefts shallow to deep, each with three stigmatic spines, median spine nearly twice as long as lateral spine; medians 60–75  $\mu\text{m}$  long, laterals 24–38  $\mu\text{m}$  long. Eyespots located near margin.

**Venter.** Derm membranous. Multilocular disc-pores 6–7  $\mu\text{m}$  wide, each with 8–10 loculi, mostly with ten loculi, abundant around vulvar area; one or two transverse rows on each abdominal segments; and also small groups present laterad of each metacoxa and mesocoxa, but not observed around procoxa. Spiracular pores 4–5  $\mu\text{m}$  wide, each with five loculi, in a narrow band 2–4 pores wide between each spiracle and stigmatic cleft. Ventral tubular ducts of three types: Type I each with 33–39  $\mu\text{m}$  long, with a developed outer ductule and a moderately broad inner ductule, straight or slightly curved, with a flower-shaped terminal gland, densely present on posterior medial area of head; frequent on anterior medial area of prothorax, extending to inner submarginal area of thorax; and also sparsely scattered on inner submarginal area of abdomen; Type II each with 16–24  $\mu\text{m}$  long, with a narrow inner ductule and a large flower-shaped terminal gland, rarely present on inner submarginal area and posterior medial area of abdomen; Type III each with 15–24  $\mu\text{m}$  long, with a long filamentous inner ductule and a quite small terminal gland, moderately present on submarginal area between anal clefts and each posterior spiracular furrow; all types of ventral tubular ducts absent on medial area of meso- and metathorax and anterior abdomen, and submarginal area of head. Ventral microducts present on entire venter, especially frequent on sub-



**Figure 5.** *Coccus malloti* (Takahashi, 1956), adult female. **A** dorsal microduct **B** dorsal tubercle **C** dermal areolations **D** dorsal seta **E** stigmatic spines **F** marginal seta **G** anal plates **H** preopercular pore **I** multilocular disc-pores **J** ventral tubular duct (Type II) **K** leg **L** ventral tubular duct (Type III) **M** ventral tubular ducts (Type I) **N** ventral seta **O** spiracular pore **P** ventral microduct **Q** antenna. Scale bars: 200  $\mu\text{m}$  (**K**, **Q**); 100  $\mu\text{m}$  (**G**); 50  $\mu\text{m}$  (**C**, **E**); others = 10  $\mu\text{m}$ .

margin. Ventral setae with three pairs of long pregenital setae; approximately three or four pairs of long and short setae between antennae; other setae sharply spinose, each 5–10  $\mu\text{m}$  long, sparsely distributed over entire venter. Legs well developed, each with a tibio-tarsal articulation and an articulatory sclerosis; total length of each metathoracic leg 614–769  $\mu\text{m}$  long; each coxa 141–189  $\mu\text{m}$  long, trochanter+femer 209–257  $\mu\text{m}$  long, tibia+tarsus 243–311  $\mu\text{m}$  long, claw 17–24  $\mu\text{m}$  long. Tarsal digitules thinner and longer than claw digitules. Spiracles normal, mostly posterior peritreme broader than anterior: anterior peritremes each 37–52  $\mu\text{m}$  wide, posterior peritremes each 45–61  $\mu\text{m}$  wide. Antenna 8-segmented, each 339–399  $\mu\text{m}$  long. Clypeolabral shield 138–157  $\mu\text{m}$  wide.

**Host plant.** Recorded from six genera in six families (García-Morales et al. 2016). In Korea, it was found on *Aphananthe aspera* (Cannabaceae) and *Ilex cornuta* (Aquifoliaceae).

**Distribution.** Only known from Japan (Takahashi 1956); first record for Korea.

**Comments.** The above description based on Korean specimens agrees well with that of Takahashi (1956), except that variation in the number of dorsal tubercles and marginal setae, and exact distributions of each type of ventral tubular ducts are newly provided in this study. *Coccus malloti* probably has intermediate morphological characters between the tribes Coccini and Pulvinariini. However, the woolly test, known as ovisac and one of the typical characters of the Pulvinariini, is not observed in the species. In addition, some slide-mounted specimens of *C. malloti* contain eggs and nymphs in their body, which indirectly indicates that they would not produce an ovisac for oviposition. Although *Coccus malloti* would be retained in the tribe Coccini, it needs to be reviewed with its type materials to clarify the exact generic position of the species.

### *Coccus pseudomagnoliarum* (Kuwana, 1914)

Fig. 6A–B

*Lecanium (Eulecanium) pseudomagnoliarum* Kuwana, 1914: 7.


**Material examined.** Adult female: Sujeong-dong, Yeosu-si, Jeollanam-do, 27.v.2015, coll. J.Y. Choi, on *Celtis* sp. (Cannabaceae), 10♀♀.

**Diagnosis.** Adult females in life (Fig. 6A–B) greenish or greyish, with light or dark yellow mottling; dermal areolations present; dorsal tubercles absent; dorsal tubular ducts absent; dorsal setae with sharply pointed apices; marginal setae with simple pointed or spatulate apices; multilocular disc-pores usually with 6–10 loculi; ventral tubular ducts with a narrow inner ductule, few present on submarginal area of posterior abdomen; antennae each 8-segmented; legs without tibio-tarsal articulatory scleroses.

**Host plant.** Recorded from six genera in five families (García-Morales et al. 2016). In Korea, it has been recorded from the following plants: *Celtis willdenowiana* (Cannabaceae), *Citrus* sp., *Phellodendron amurense*, *Poncirus trifoliata* (Rutaceae), *Clerodendron trichotomum* (Lamiaceae), and *Zelkova serrata* (Ulmaceae) (Paik 1978).

**Distribution.** Mainly known from Palearctic Region including Australia, Europe, Iran, Israel, Japan, Russia, Korea, and USA (García-Morales et al. 2016).





**Figure 6.** *Coccus pseudomagnoliarum* (Kuwana, 1914). **A** immature adult females **B** mature adult female and 1<sup>st</sup> instars.

## Discussion

Recently, Lin et al. (2017) described *Coccus praetermissus* Lin & Tanaka, which could be confused with a cosmopolitan species, *Coccus hesperidum* Linnaeus, 1758, based on morphological and molecular evidences. They pointed out that a morphological difference exists between the adult females of two genetically distinct species, the shape of dorsal setae, although molecular data (COI) should be used for exact identification. The adult female of *Coccus praetermissus* has dorsal setae with bluntly rounded apices, whereas those of *C. hesperidum* have sharply pointed tips. *Coccus ficicola* sp. n. is close to *C. praetermissus* in having the former type of dorsal setae, but differs in the distributional pattern of ventral tubular ducts, which is a reliable and constant character in each species within the genus *Coccus*. The ventral tubular ducts of *Coccus ficicola* are present on medial area of head, pro- and mesothorax, and submarginal area of abdomen, whereas *C. praetermissus* has the structures on medial area of mesothorax only.

Under the morphological comparison with congeners, we conclude that *Coccus ficicola* sp. n. is a distinct species which is a morphologically differentiated lineage. The distinctive pattern of ventral tubular ducts seems to be an autapomorphic feature of *Coccus ficicola* because it shows uniqueness in the comparison of morphological characters. In order to clarify the phylogenetic relationships of a new species within the genus *Coccus*, molecular analysis employing mitochondrial and nuclear loci are required.

## Acknowledgments

We thank Dr. Sora Kim for reviewing the manuscript for this study. We are also grateful for a colleague, Mr. Seunghyun Lee (Department of Agricultural Biotechnology, Seoul National University, Seoul, Korea), who provided specimens. We also thank Dr. Chris J. Hodgson (Department of Biodiversity and Biological Systematics, The National Museum of Wales, Cardiff, UK), Dr. T. Kondo (Corporación Colombiana

de Investigación Agropecuaria, Corpoica, Colombia) and Dr. Yen P. Lin (School of Biological Sciences, The University of Queensland, Brisbane, Australia) for the useful comments. In addition, we thank Dr. H. Tanaka (Ehime University, Ehime, Japan) for comparing our samples with Japanese samples.

The research on the species newly recorded in Korea was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR201701203).

## References

- Avasthi RK, Shafee SA (1991) Revision of the genus *Coccus* Linn. in India (Insecta, Homoptera, Coccidae). *Journal of the Bombay Natural History Society* 88(3): 329–348.
- Ben-Dov Y (1981) A new species of *Coccus* (Homoptera: Coccidae) from mango in Israel, and a re-description of *C. gymnospori* (Green). *Bulletin of Entomological Research* 71(04): 649–654. <https://doi.org/10.1017/S0007485300010178>
- Danzig EM, Gavrillov-Zimin IA (2014) Palaearctic mealybugs (Homoptera: Coccinea: Pseudococcidae). *Fauna of Russia and Neighboring Countries (new series)* 148, 678 pp.
- De Lotto G (1957) On some Ethiopian species of the genus *Coccus* (Homoptera: Coccoidea: Coccidae). *Journal of the Entomological Society of southern Africa* 20: 295–314.
- De Lotto G (1959) Further notes on Ethiopian species of the genus *Coccus* (Homoptera: Coccoidea: Coccidae). *Journal of the Entomological Society of southern Africa* 22: 150–173.
- De Lotto G (1960) The green scales of coffee in Africa south of the Sahara (Homoptera, Coccidae). *Bulletin of Entomological Research* 51: 389–403. <https://doi.org/10.1017/S0007485300058041>
- García-Morales M, Denno BD, Miller DR, Miller GL, Ben-Dov Y, Hardy NB (2016) ScaleNet: A literature-based model of scale insect biology and systematics. <http://scalenet.info>, <https://doi.org/10.1093/database/bav118>
- Gavrillov-Zimin IA (2013) New scale insects (Homoptera: Coccinea) from Sulawesi and New Guinea, with some other additions to the Indonesian fauna. *Tropical Zoology* 26: 64–86. <https://doi.org/10.1080/03946975.2013.807570>
- Gill RJ, Nakahara S, Williams ML (1977) A review of the genus *Coccus* Linnaeus in America north of Panama (Homoptera: Coccoidea: Coccidae). *Occasional Papers in Entomology, State of California Department of Food and Agriculture* 24: 1–44.
- Granara de Willink MC, Pirovani VD, Ferreira PS (2010) Las Especies de *Coccus* que Afectan *Coffea arabica* en Brasil (Coccoidea: Coccidae) y Redescripción de Dos Especies. [*Coccus* Species affecting *Coffea arabica* in Brazil (Coccoidea: Coccidae) and the redescription of two Species.]. *Neotropical Entomology* 39(3): 391–399. <https://doi.org/10.1590/S1519-566X2010000300013>
- Green EE (1896) Catalogue of Coccidae collected in Ceylon. *Indian Museum Notes* 4: 2–10.
- Green EE (1904) The Coccidae of Ceylon. Pt. III. Dulau & Co., London, UK, 171–249.
- Green EE (1908) Remarks on Indian scale insects (Coccidae), Part III. With a catalogue of all species hitherto recorded from the Indian continent. *Memoirs of the Department of Agriculture in India, Entomology Series* 2: 15–46.



- Green EE (1922) Supplementary notes on the Coccidae of Ceylon. Part IV. Journal of the Bombay Natural History Society 28: 1007–1037.
- Hodgson CJ, Henderson RC (2000) Coccidae (Insecta: Hemiptera: Coccoidea). Manaaki Whenua Press, Lincoln, Canterbury, New Zealand, 259 pp.
- Hodgson CJ (1994) The scale insect family Coccidae: an identification manual to genera. CAB International Wallingford, Oxon, UK, 639 pp.
- Kapranas A, Morse JG, Pacheco P, Forster LD, Luck RF (2007) Survey of brown soft scale *Coccus hesperidum* L. parasitoids in southern California citrus. Biological Control 42(3): 288–299. <https://doi.org/10.1016/j.biocontrol.2007.05.012>
- Kondo T, Hardy N (2008) Synonymy of *Etiennaea* Matile-Ferrero with *Hemilecanium* Newstead (Hemiptera: Coccidae), based on morphology of adult females, adult males and first-instar nymphs, and description of a new potential pest species from the Ryukyu Archipelago, Japan. Entomological Science 11: 189–213. <https://doi.org/10.1111/j.1479-8298.2008.00268.x>
- Kuwana SI (1914) Coccidae of Japan, V. Journal of Entomology and Zoology 6: 1–8.
- Lin YP, Kondo T, Gullan P, Cook LG (2013) Delimiting genera of scale insects: molecular and morphological evidence for synonymising *Taiwansaissetia* Tao, Wong and Chang with *Coccus* Linnaeus (Hemiptera: Coccoidea: Coccidae) Systematic Entomology 38(2): 249–264. <https://doi.org/10.1111/j.1365-3113.2012.00664.x>
- Lin YP, Tanaka H, Cook LG (2017) A newly recognised species that has been confused with the global polyphagous pest scale insect, *Coccus hesperidum* Linnaeus (Hemiptera: Coccoidea: Coccidae). Zootaxa 4320(3): 571–591. <https://doi.org/10.11646/zootaxa.4324.3.10>
- Linnaeus C (1758) Systema Naturae, per Classes, Ordines, Genera, Species cum Characteribus, Differentiis, Synonymis, Locis. Ed X, Triformata, Tomus I. Laurentii Salvii, Holmiae, 823 pp.
- Newstead R (1898) Observations on Coccidae (No. 17). Entomologist's Monthly Magazine 34: 92–99.
- Opinion 1303 (1985) *Coccus* Linnaeus, 1758 and *Parthenolecanium* Sulc, 1908 (Insecta, Hemiptera, Homoptera): type species designated. Bulletin of Zoological Nomenclature 42: 139–141.
- Paik WH (1978) Illustrated flora and fauna of Korea. Insecta (VI). No. 22. Min. Education, 481 pp.
- Starr F, Starr K, Loope L (2003) *Ficus benghalensis*. Datasheet, Hawaiian Ecosystems at Risk project. [www.cabi.org/isc/datasheet/24066](http://www.cabi.org/isc/datasheet/24066)
- Takahashi R (1956) *Pulvinaria* of Japan (Coccidae, Homoptera). Kontyu 24: 23–30.
- Tao CCC, Wong CY, Chang YC (1983) Monograph of Coccidae of Taiwan, Republic of China (Homoptera: Coccoidea). Journal of Taiwan Museum 36(1): 57–107.
- Waller JM, Bigger M, Hillocks RJ (2007) Coffee pests, diseases and their management. CAB International Wallingford, Oxon, UK, 434 pp. <https://doi.org/10.1079/9781845931292.0000>
- Wang F, Feng JN (2012) A new species of *Coccus* (Hemiptera, Coccoidea, Coccidae) from China. ZooKeys 244: 59–65. <https://doi.org/10.3897/zookeys.244.4045>
- Williams DJ, Watson GW (1990) The scale insects of the tropical South Pacific Region. Pt. 3: The soft scales (Coccidae) and other families. CAB International, Wallingford, Oxon, 267 pp.