

INSECTA MUNDI

A Journal of World Insect Systematics

0347

Updated list of *Pinnaspis* armored scales (Hemiptera: Diaspididae)
of Korea

Soo-Jung Suh
Animal and Plant Quarantine and Inspection Agency
476 Dongtanjiseongro
Suwon
South Korea

Date of Issue: February 7, 2014

Soo-Jung Suh

Updated list of *Pinnaspis* armored scales (Hemiptera: Diaspididae) of Korea
Insecta Mundi 0347: 1–9

ZooBank Registered: urn:lsid:zoobank.org:pub:3F9D4643-F0C3-47F7-BC03-03E8D56C08B4

Published in 2014 by

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Eugenio H. Nearn, e-mail: gino@nearns.com

Production editors: Michael C. Thomas, Paul E. Skelley, Brian Armitage, Ian Stocks, Eugenio H. Nearn

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, Joe Eger, A. Rasmussen, Gary Steck, Ian Stocks, A. Van Pelt, Jennifer M. Zaspel, Nathan P. Lord, Adam Brunke

Spanish editors: Julieta Brambila, Angélico Asenjo

Website coordinator: Eugenio H. Nearn

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, Great Britain

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Author instructions available on the Insecta Mundi page at:

<http://centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Updated list of *Pinnaspis* armored scales (Hemiptera: Diaspididae) of Korea

Soo-Jung Suh

Animal and Plant Quarantine and Inspection Agency

476 Dongtanjiseongro

Suwon

South Korea

suhSJ97@gmail.com

Abstract. *Pinnaspis chamaecyparidis* Takagi, *Pinnaspis hikosana* Takagi and *Pinnaspis uniloba* (Kuwana), occurring on *Chamaecyparis obtusa* Endl., *Styrax japonica* S. et Z. and *Cleyera japonica* Thunb. are newly documented in the Korean fauna of armored scales (Diaspididae). The characters of these species are here redescribed with illustrative photographs and information on distribution and hosts along with a dichotomous key to the species of *Pinnaspis* for correct species identification. In addition, the paper discusses the current status of *Pinnaspis buxi* (Bouché) and *Pinnaspis strachani* (Cooley) which are known as native armored scale insects of Korea by analyzing information on the result of the survey.

Key words. *Pinnaspis chamaecyparidis*, *Pinnaspis hikosana*, *Pinnaspis uniloba*, New Record, Diaspididae, Korea

Introduction

Species in the genus *Pinnaspis* Cockerell (Hemiptera: Diaspididae) have a more or less slender, fusiform and elongate body, the metathorax distinctly or slightly protruding lateral lobes and the pygidium triangular or roundish. The most distinctive feature is the pair of median lobes, which are united at the base by an elongate zygois; the inner edges of the median lobes are parallel and very close together at the apices, appearing to be fused (Takagi 1970; Williams and Watson 1988). According to the ScaleNet database, 46 species have been documented around the world (Ben-Dov et al. 2013); of these, 41 species and 12 species were reported to be originated from Oriental (89%) and Palaearctic (26%) regions, respectively. Between 1963 and 2003, Takagi discovered and described 14 *Pinnaspis* species from eastern Asia, thus this illustrates the abundance and diversity of the genus in the eastern Asia (Ben-Dov et al. 2013).

Species of the genus *Pinnaspis* species were first time recorded in Korea by Nakayama and Kanda in 1933 and 1942 (cited from the publication of Paik (2000) who documented the presence of *Pinnaspis aspidistrae* (Signoret), *Pinnaspis buxi* (Bouché) and *Pinnaspis strachani* (Cooley)). Since that time, no species have been added to the Korean armored scale fauna (Paik 1978; Paik 2000; Lee 2010), although some species of the genus have been found in other eastern Asian countries including Japan and China. As the result of the Korean Diaspididae project survey initiated in 2006, the following three additional *Pinnaspis* species occurring on *Chamaecyparis obtusa* Endl. (Cupressaceae), *Styrax japonica* S. et Z. (Styracaceae) and *Cleyera japonica* Thunb. (Theaceae), respectively were found: *Pinnaspis chamaecyparidis* Takagi, *Pinnaspis hikosana* Takagi and *Pinnaspis uniloba* (Kuwana) and represent the first records of the occurrence of these species in Korea. All three species were collected on leaves and twigs and no damage to their host plants was observed.

Regarding the economic importance of *Pinnaspis*, *P. aspidistrae* and *P. strachani* are known for polyphagous and worldwide pests. The fern scale, *P. aspidistrae* is probably native to the Oriental region (Miller and Davidson 2005) and is considered to be one of the principal armored scale pests of the world (Beardsley and González 1975). Also the fern scale is regarded as serious pest of fern, foliage, citrus, oil palm and rubber trees in the USA, Brazil, Chile, Japan and Malaysia (Miller and Davidson 2005). In Korea, it was reported to occur only on *Asplenium nidus* (Aspleniaceae), *Aspidistra elatior* (Asparagaceae) and undetermined Orchidaceae in greenhouses (Paik 1978). During the recent survey, it was often collected not only in greenhouses but also on outside plants. Based on the material in the PQTC's collection, it was found on *Liriope platyphylla* (Asparagaceae) and *Rhaphis humilis* (Arecaceae) outdoors in Korea. The lesser snow scale, *P. strachani*, described from Nigeria, is a serious and wide-

spread pest as well. It occasionally causes serious damage to hibiscus, tamarind, citrus, black pepper, arecanut, olives and several wild plants (Miller and Davidson 2005).

While updating the list of the *Pinnaspis* armored scales of Korea, the presence of *P. buxi* and *P. strachani* in Korea could not be confirmed because no specimens could be found from collections made from 2006 to the present time, although according to the literature (Paik 1978; Paik 2000; Lee 2010), both species have been found in greenhouses in Korea.

The purpose of this paper is: 1) to update the list of the *Pinnaspis* species that occur in Korea, 2) to provide an identification key to species of *Pinnaspis* in Korea, 3) to provide photos, redescriptions for the three species newly documented in Korea, and information on the plant hosts and distribution of species, and 4) to review the current status of *P. buxi* and *P. strachani* in Korea.

Materials and Methods

All slide-mounted specimens used for this paper are deposited in the Collection of Plant Quarantine Technology Center (PQTC), QIA (Figure 3F). The dichotomous key mentioned below was based on specimens of adult females collected on leaves and twigs of their plant hosts. Terminology for morphological structures used in descriptions and an identification key follows that of Miller and Davidson (2005). Photographs were taken using an AxioCam MRc5 camera through ZEISS Axio Imager M2 Microscope and a Leica M165C microscope with Delta pix camera. An asterisk(*) is used to indicate a new distribution record. The following *Pinnaspis* species: *P. aspidistrae*, *P. buxi* and *P. strachani* were well described in the book of Paik (1978), and are not addressed here.

Description

Pinnaspis chamaecyparidis Takagi, 1961

(Figures 1A–F)

Pinnaspis chamaecyparidis Takagi, 1961: 72–73. Type data: Japan, on *Chamaecyparis obtusa*, 8-i-1955 and 16-iv-1956.

Field Characters. Adult female cover narrowly to broadly oyster-shell shaped, light to dark brown; shed skins margin, yellow brown to brown. Male cover smaller, felted, white, elongate, with slight median carina; shed skin yellowish.

Slide-mounted Characters. Adult female with 2 pairs of well developed lobes, third lobes inconspicuous; paraphyses attached to medial margin of median lobules of second lobe. Median lobes small in size, closely appressed together, slightly elongate with 1–2 deeply lateral notches, the basal zygonis robust, and distinctly produced anteriorly into apex of pygidium; second lobes with 2 lobules, medial lobules slightly expanded apically, with a pair of slender basal sclerites. Gland spine formula 1–1–1, with about 14 gland spines near each body margin from segments 5 to 7. Macroducts of 2 sizes, larger size in marginal areas on segments 4 to 7, in submarginal areas on segments 1 to 3, smaller size in submarginal and marginal areas from metathorax to segment 3. Perivulvar pores in 5 groups, 38–48 pores. Perispiracular pores usually with 3 loculi, anterior spiracles each with 3–6 pores, posterior spiracles with 1–2 pores. Anal opening located about middle of pygidium. Antennae each with 1 conspicuous seta. Body elongated fusiform with prepygidial segments each slightly convex laterally. Preanal scars absent.

Affinities. This species is most similar to *Pinnaspis juniper* Takahashi which is not known to occur in Korea, from which it differs mainly by the second lobes which are well developed, with the inner lobule slightly expanded apically.

Material examined. Korea. Gyeongsangnamdo (GN): Dadaepo-dong, Saha-gu, Busan-si, 5 adult females, on *Chamaecyparis obtusa* (Cupressaceae) leaf, 13-xii-2007 (S.J. Suh); Dongbaekseom, 2 adult




Figure 1. *Pinnaspis chamaecyparidis* Takagi. A) Habitus. B) Female. C) Antennae. D) Anterior and posterior spiracles. E) Median lobes. F) Perivulvar pores.

females, on *Juniperus chinensis* (Cupressaceae) leaf, 14-viii-2008 (S.J. Suh); Dongbaek park, 3 adult females and 2 nymphs, on *Juniperus chinensis* (Cupressaceae) leaf, 24-ix-2008 (S.J. Suh); Tongdosa, 7 adult females, on *Chamaecyparis obtusa* (Cupressaceae) leaf, 8-ix-2009 (S.J. Suh). Jeollabukdo (JB): Gyeongwon-dong, Wansan-gu, Jeonju-si, 5 adult females, on *Chamaecyparis obtusa* (Cupressaceae) leaf, 17-iii-2006 (S.J. Suh). Jeollanamdo (JN): Gatbawi, 7 adult females, on *Juniperus chinensis* (Cupressaceae) leaf, 23-vii-2012 (S.J. Suh).

Hosts. This species has a restricted host range occurring on conifers. Cupressaceae: *Chamaecyparis obtusa*, *Chamaecyparis pisifera*, *Cryptomeria japonica*, *Juniperus chinensis* (Kawai 1980; Ben-Dov et al. 2013).

Distribution. Japan and Korea* (Ben-Dov et al. 2013).

Remarks. This species has two generations per year and hibernates as an adult female in Japan (Kawai 1980).

***Pinnaspis hikosana* Takagi, 1961**
(Figures 2A–F)

Pinnaspis hikosana Takagi, 1961: 73–74. Type data: Japan, on *Viburnum* sp., 10-v-1957.

Field Characters. Adult female cover narrowly to broadly oyster-shell shaped, dark brown; shed skins marginal, yellow brown to brown. Male cover smaller, felted, white, elongate, with tricarinate (Kawai 1980).

Slide-mounted Characters. Adult female with 2 pairs of lobes, third lobes inconspicuous; paraphyses usually absent. Median lobes prominent, large in size, closely appressed together, a half circle shaped with 1–2 deeply lateral notches, the basal zygonis small, and slightly produced anteriorly into apex of pygidium; second lobes present and represented by a small, apically pointed process and lateral lobules by a small inconspicuous process. Gland spine formula 1–1–1, with about 16–18 gland spines near each body margin from segments 2 to 7. Macroducts of 2 sizes, larger size in marginal areas on segments 4 to 7, smaller size in submarginal and marginal areas from metathorax to segment 3. Perivulvar pores in 5 groups, 48–58 pores. Perispiracular pores usually with 3 loculi, anterior spiracles each with 5–6 pores, posterior spiracles with 1–2 pores. Anal opening located about middle of pygidium. Antennae each with 1 conspicuous seta. Body elongated fusiform with prepygidial segments each moderately convex laterally. Preanal scar present and represented by slender crescent shaped scars.

Affinities. This species is very similar to *P. strachani* from which it differs mainly by the basal zygonis of medial lobes which are small and slightly produced anteriorly into apex of pygidium and the second lobes present and represented by a small, apically pointed process.

Material examined. Korea. Chungcheongnamdo (CN): Gyeryosan, 7 adult females, on *Styrax japonica* (Styracaceae) twig, 26-iv-2013 (S.J. Suh).

Hosts. Anacardiaceae: *Lannea coromandelica*. Aquifoliaceae: *Ilex crenata*, *Ilex integra*. Caprifoliaceae: *Viburnum dilatatum*, *Viburnum plicatum*, *Viburnum sieboldii*, *Viburnum* sp. Cornaceae: *Cornus controversa*, *Cornus macrophylla*, *Cornus* sp. Clethraceae: *Clethra barbinervis*. Daphniphyllaceae: *Daphniphyllum macropodum*. Styracaceae: *Styrax japonica*. Theaceae: *Camellia japonica*, *Cleyera japonica*, *Eurya japonica* (Kawai 1980; Ben-Dov et al. 2013).

Distribution. Japan and Korea* (Ben-Dov et al. 2013).




Figure 2. *Pinnaspis hikosana* Takagi. A) Habitus. B) Female. C) Antennae. D) Anterior and posterior spiracles. E) Median lobes and preanal scars. F) Perivulvar pores.

Remarks. This species has two generations per year, hibernates as an adult female and the first generation appears in mid-May to early June in Japan (Kawai 1980).

***Pinnaspis uniloba* (Kuwana), 1909** (Figures 3A–E)

Mytilaspis (*Lepidosaphes*) *uniloba* Kuwana, 1909: 156. Type data: Japan, on *Osmanthus* sp., ?-iii-1907.

Field Characters. Adult female cover elongate oyster-shell shaped, dark brown (Takagi 1970). Male not observed during the survey.

Slide-mounted Characters. Adult female with one lobe, the second lobes absent. Median lobes fused throughout their entire length, thus forming a single median lobe with 2–3 lateral notches, the basis zygosis prominent, small and slightly produced anteriorly into apex of pygidium, the basal scleroses heavily sclerotized patches. Gland spine formula 1–1–1, with about 16–18 gland spines near each body margin from segments 3 to 7. Macroducts of 2 sizes, larger size in marginal areas on segments 4 to 7, smaller size in marginal areas from metathorax to segment 3 and in submarginal area of segment 4. Perivulvar pores in 5 groups, 42–60 pores. Perispiracular pores usually with 3 loculi, anterior spiracles each with 2–4 pores, posterior spiracles nothing or 1 pore. Anal opening located about middle of pygidium. Antennae each with 1 conspicuous seta. Body elongated fusiform, almost parallel on the lateral margins through the thorax and prepygidial abdominal segments, with prepygidial segments each slightly convex laterally, with pygidium triangular. Preanal and poster lateral to the anus scars present and represented by a slender sclerotized bar.

Affinities. This species is distinct among species of *Pinnaspis* that occur in Korea by having a wholly fused median lobe.

Material examined. Korea. Jeollanamdo (JN): Wando arboretum, 13 adult females, on *Cleyera japonica* (Theaceae) leaf, 10-v-2007 (S.J. Suh).

Hosts. Apocynaceae: *Alyxia olivaeformis*, *Alyxia* sp. Fabaceae: *Bauhinia racemosa*. Magnoliaceae: *Michelia* sp. Oleaceae: *Osmanthus fortunei*, *Osmanthus fragrans*, *Osmanthus* sp. Rutaceae: *Aegle marmelos*. Theaceae: *Adinandra milletti*, *Adinandra ryukyuensis*, *Adinandra* sp., *Camellia japonica*, *Cleyera japonica*, *Cleyera ochracea*, *Eurya japonica*, *Thea japonica* (Kawai 1980; Ben-Dov et al. 2013).

Distribution. Hawaiian Islands, India, Taiwan, China, Japan and Korea* (Ben-Dov et al. 2013).

Remarks. This species has one generation per year, hibernates in the egg stage and the first generation appears in May in Japan. It lacks males and is parthenogenetic (Kawai 1980).

Key to species of *Pinnaspis* from Korea (slide mounted adult female)

1. Found only on conifers. ***P. chamaecyparidis* Takagi**
- Not found on conifers. **2**
2. Median lobes fused throughout their length, thus forming a single median lobe. ***P. uniloba* (Kuwana)**
- Median lobes although usually very closely appressed together, separate for at least some distance from the apex, if not to the base. **3**
3. Median lobes protrude less than or about the same distance as second lobes. **4**
- Median lobes protrude beyond or about the same distance as second lobes. **5**
4. With submarginal ducts on dorsum of 5th abdominal segment. ***P. aspidistrae* (Signoret)**
- Without submarginal ducts on dorsum of 5th abdominal segment. ***P. buxi* (Bouché)**
5. Basal zygosis of median lobes slender, scarcely produced anteriorly beyond bases of the lobes. ***P. hikosana* Takagi**
- Basal zygosis of median lobes robust, distinctly produced anteriorly beyond bases of the lobes. ***P. strachani* (Cooley)**




Figure 3. Three species of *Pinnaspis*. **A)** *P. uniloba* (Kuwana), female. **B)** *P. uniloba* (Kuwana), antennae. **C)** *P. uniloba* (Kuwana), anterior and posterior spiracles. **D)** *P. uniloba* (Kuwana), median lobes and preanal scars. **E)** *P. uniloba* (Kuwana), perivulvar pores. **F)** Slide-mounted specimens.

Discussion

The list of Korean *Pinnaspis*, including *Pinnaspis chamaecyparidis* Takagi, *Pinnaspis hikosana* Takagi and *Pinnaspis uniloba* (Kuwana) which are newly recorded, is updated through the survey conducted recently. The information provided in the paper will be helpful in better understanding Korean *Pinnaspis* species. Such research should be considered to be an integral part of more appropriate and cost-effective preventative quarantine procedures to overcome the challenge from exotic species.

Of these, *P. buxi* and *P. strachani* which have been reported to occur in Korea should be reviewed by analyzing information on their morphological affinities and the results of the survey. First, *P. buxi* is very near *P. aspidistrae* and can only be separated on the basis of the laterally produced metathoracic lobes of *P. aspidistrae* and fewer submarginal ducts of *P. buxi*. It is likely that this record represents a misidentification of *P. aspidistrae*. The other species, *P. strachani* is most similar to *P. aspidistrae*; a few specimens are difficult or impossible to determine using the criteria, the combination of the following characters; the fern scale has small median lobes that protrude less than or about the same distance as the large second lobes, the preanal scars are lacking or represented by light sclerotized patches and the pores around posterior spiracles are present; the lesser snow scale has large median lobes that protrude beyond or about the same distance as the small second lobes, the preanal scars are represented by heavily sclerotized bars and pores around posterior spiracles are absent (Ferris and Rao 1947; Takagi 1963, 1970; Miller and Davidson 2005). The resolution of the taxonomy of the species comprising the *aspidistrae-strachani* complex of *Pinnaspis* has long remained unsettled. *Pinnaspis aspidistrae* has sometimes been misidentified as *P. strachani* and *vice-versa*. Although the two species were recorded in Korea, they are considered to have failed to establish in the exterior environment and greenhouses because they have not been collected in the field and greenhouses during the last 7 years. Therefore, the presence of both of these species in Korea should be regarded as uncertain.

Acknowledgements

I would like to thank Dr. Sadao Takagi (Hokkaido University, Japan) for confirming the identification. I am also grateful to Greg Evans (USDA/APHIS/NIS, Washington, DC, USA) and Greg Hodges (FDACSDPI, Florida, USA) for their useful editorial contributions to early drafts of this paper.

Literature Cited

- Ben-Dov, Y., D.R. Miller, and G.A.P. Gibson. 2013. ScaleNet(webpage) [http:// www.sel. barc.usda. gov/scalenet/scalenet.htm](http://www.sel.barc.usda.gov/scalenet/scalenet.htm). Accessed October 2013.
- Beardsley, J.W., and R. H. González. 1975. The biology and ecology of armored scales. Annual Review of Entomology 20: 47–73.
- Ferris, G.F., and V.P. Rao. 1947. Contribution No. 54. The genus *Pinnaspis* Cockerell (Homoptera: Coccoidea: Diaspididae). Microentomology 12: 25–58.
- Kanda, S. 1942. Studies on Coccidae from Corea (III). Insect world 46(2): 35–42.
- Kawai, S. 1980. Scale insects of Japan in colors. National Agricultural Education Association; Tokyo. 455 p.
- Kuwana, S.I. 1909. Coccidae of Japan (III). First supplemental list of Japanese Coccidae or scale insects with description of eight new species. Journal New York Entomological Society 17: 150–164.
- Lee, Y.J. 2010. Family Diaspididae. p. 80–81. In: M.K. Paek (ed.), Checklist of Korean insects. Nature and Ecology; Seoul. 598 p.
- Miller, D.R., and J.A. Davidson. 2005. Armored scales insects pests of trees and shrubs (Homoptera: Diaspididae), Cornell University Press; Ithaca, NY. 442 p.
- Nakayama, S. 1933. Notes on scale insects and host plants from Korea. Oyo-Dobutsugaku-Zasshi 3: 226–229.
- Paik, J.C. 2000. Economic Insects of Korea 6, Homoptera (Coccinea), Insecta Koreana Suppl. 13. National Institute of Agricultural Science and Technology; Seoul. 193 p.

- Paik, W.H. 1978.** Illustrated flora and fauna of Korea, vol. 22, Insecta (VI), Coccoidea. Samhwa Publishing Company; Seoul. 481 p.
- Takagi, S. 1961.** A contribution to the knowledge of the Diaspididini of Japan (Homoptera: Coccoidea) Pt. II. Insecta Matsumurana 24: 4–42.
- Takagi, S. 1963.** Notes on the Aspidistrae-complex of *Pinnaspis* with description of a new species (Homoptera: Coccoidea). Insecta Matsumurana 26(1): 64–68.
- Takagi, S. 1970.** Diaspididae of Taiwan based on material collected in connection with the Japan-U.S. cooperative science programme, 1965 (Homoptera: Coccoidea) Pt. II. Insecta Matsumurana 33: 1–146.
- Williams, D.J., and G.W. Watson.** 1988. The scale insects of the tropical south pacific region: Pt. 1, The armoured scales (Diaspididae). CAB International Institute of Entomology; London. 290 p.

Received January 3, 2014; Accepted January 30, 2014.

