

A new species of *Pseudaulacaspis* MacGillivray, 1921 from China (Hemiptera, Coccoidea, Diaspididae) with a key to Chinese species

Jiu-Feng Wei^{1,2,†}, Ji-Nian Feng^{2,‡}

1 College of Life Sciences, Northwest A & F University **2** Key Laboratory of Plant Protection Resources and Pest Management, Ministry of Education, Entomological Museum, Northwest A & F University, Yangling, Shaanxi Province, 712100, China

† urn:lsid:zoobank.org:author:E6F808B7-EC44-4934-8E37-2A8D83FD9BE2

‡ urn:lsid:zoobank.org:author:16913882-73AC-47FE-9C89-5B61FC961524

Corresponding author: Ji-Nian Feng (jinianf@nwsuaf.edu.cn)

Academic editor: Mike Wilson | Received 25 March 2012 | Accepted 9 July 2012 | Published 24 July 2012

urn:lsid:zoobank.org:pub:C91C9813-E74E-4B11-A44A-BFE31AF03706

Citation: Wei J-F, Feng J-N (2012) A new species of *Pseudaulacaspis* MacGillivray, 1921 from China (Hemiptera, Coccoidea, Diaspididae) with a key to Chinese species. ZooKeys 210: 9–17. doi: 10.3897/zookeys.210.3122

Abstract

A new species of armored scale, *Pseudaulacaspis zhenyuanensis* Wei & Feng, **sp. n.** is described and illustrated from specimens collected on *Spermatidictyon suaveolens* in China. A key to armored scale species known from China is provided.

Keywords

Hemiptera, armored scale, taxonomy, Diaspididae, new species

Introduction

The Coccoidea is one of the four superfamilies of the monophyletic suborder Setenorrhyncha belongs to the Hemiptera (Gullan and Cook 2007), with at least 30 families and around 8000 species (Andersen 2010). The family Diaspididae is the largest family of the Coccoidea with more than 2400 diaspidid species currently known (Ben-dov 2012). The higher classification within the family is uncertain but two of the major

subfamilies are the Aspidiotinae and the Diaspidinae, and most species can be assigned to one or the other (Miller and Davidson 2005)

The genus *Pseudaulacaspis* was established by MacGillivray (1921) for *Diaspis pentagona* Targioni Tozzetti, 1886 which belongs to subfamily Diaspidinae. When he described it, he referred to it 9 nominal species, which are now considered to represent only 2 species. Since then, many additional species were described and added to *Pseudaulacaspis* by other authors (Chen 1983; Ferris 1953, 1955; Hu 1986; Takagi 1956, 1961, 1966, 1970, 1985; Talng 1986, 1988; Williams and Watson 1988; Hodgson and Lagowska 2011). This genus is large with 68 species (Hodgson and Lagowska 2011) which is a widespread and polyphagous genus infesting a large number of plant (Borchsenius 1966) and occurs in most of zoogeographical regions except Antarctica. Up until now, 32 species have been described from China.

In the present paper, a new species *P. zhenyuanensis* sp. n. is described and illustrated, bringing the number of recognized species in this genus to 69, of which 33 are recorded from China. And a key to species from China is included.

Materials and methods

The morphological terms for Diaspididae follow those of Henderson (2011). The illustrations of the adult female are drawn from slide-mounted specimens, which depict the dorsum on the left and venter on the right. Enlargements of important characters are shown around the edges of the main illustration. All measurements are given in micrometers (μm). Measurements were made using the measurement tools NIT-Elements D. The abbreviations L1, L2, L3 and L4 stand for median and second to fourth pygidial lobes.

All specimens are deposited in the Entomological Museum, Northwest A & F University, Yangling, Shaanxi, China (NWAUFU).

Taxonomy

Genus *Pseudaulacaspis* MacGillivray, 1921

<http://species-id.net/wiki/Pseudaulacaspis>

Pseudaulacaspis MacGillivray, 1921: 305. Type species: *Diaspis pentagona* Targioni Tozzetti, by original designation.

Generic diagnosis. Female scale. White, suborbicular or long pyriform. Exuviae terminal. **Male scale.** Same colour as female scale, elongate.

Adult female. Body shape varied, fusiform, olivary or elongate; derm membranous except for the marginal of pygidium; mesothorax, metathorax, and abdominal segments I–III produced laterally. **Cephalothorax.** Antennae each with a seta. Anterior spiracles each usually with a cluster of trilocular pores, posterior spiracle each

associate with or without trilocular pores. **Pygidium.** With 2 or 3 pairs of lobes. Median lobes (L1) well-developed, much larger than lobules of lateral lobes, zygotic basally, with a distinct pair of marginal setae between lobes. In general, L1 divide into two types: bark-type, individuals occur on bark and prominent median lobes; leaf-type, those on leaves and sunken into the pygidium. Second lobes (L2) much smaller than the L1, bilobed, divided into inner lobule and outer lobule, outer lobule usually smaller than inner, in some species much reduced. Third lobes (L3) smaller than L2, bilobed or represented by serrations along the body margin in some species. **Gland spines.** Gland spines developed, usually single on abdominal segments VI–VIII, becoming shorter into conical on anterior segments which called gland tubercles. **Ducts.** Dorsum with 2-barred ducts, forming submedial and submarginal rows on abdominal and pygidium, usually as same size as marginal macroducts. Ventral microducts scattered. **Anal opening.** Anal opening close to the base of or situated about the centre of the pygidium. **Perivulvar pores** quinquelocular, in five groups.

Remarks. This genus is very closely related to *Chionaspis* Signoret, 1868 and *Aulacaspis* Cockerell, 1893 in feature of pygidial lobes and dorsal ducts present on pygidium and abdomen, but can differ from these genus: presence of a pair of setae between the median lobes in *Pseudaulacaspis*, but absent in *Chionaspis* and *Aulacaspis*.

***Pseudaulacaspis zhenyuanensis* Wei & Feng, sp. n.**

urn:lsid:zoobank.org:act:AB75F5D8-7BDF-42D4-9DA9-13F4DFD1D43F


http://species-id.net/wiki/Pseudaulacaspis_zhenyuanensis

Figures 1–8

Material examined. Holotype: adult female: CHINA: Guizhou Prov., Zhenyuan County, 13. viii. 1996, Zeng (NWAFU).

Paratypes: 2 adult females: same data as the holotype (NWAFU).

Description, n=3. Adult female. Appearance in life not recorded. Slide-mounted adult female 1755–1910 μm long (holotype 1910 μm long); 930–970 μm wide (holotype 931 μm wide), body outline fusiform, derm membranous except for pygidium. Normally widest at metathorax and abdominal segment I, lateral abdominal lobes well-developed, with large gland spines on the margin of prepygidial and pygidial segments. **Cephalothorax.** Antennae each with 1 long fleshy seta, distance between antennae is 111 μm . Anterior spiracle each with 12–31 trilocular pores in a cluster, posterior spiracle each with 11–17 trilocular pores. **Pygidial Lobes.** With 3 pairs of lobes; L1 well-developed, zygotic basally, protruding from pygidial margin, with small serrations along both margins, with a pair of setae between lobes; L2 bilobate, inner lobule rounded, much larger than outer lobule; L3 bilobate, slightly smaller than L2, inner lobule rounded, outer lobule margin serrate; L4 represented by serrations along the body margin. **Gland spines.** Large, arranged singly on pygidial segments VI–VIII but with 2 on segment V, 4–5 on segment IV, 5–6 on segment III, 5 on segment II, anterior spines smallest (on seg-



Figures 1–8. *Pseudaulacaspis zhenyuanensis* Wei & Feng, sp. n., adult female: **1** habitus **2** antennae **3** anterior spiracle **4** gland tubercles **5** detail of gland macroduct **6** detail of the duct in the head on the dorsum **7** detail of pygidium **8** pygidium.

ment II). **Gland tubercle** present submarginally, with 2 on prothorax, 9-11 on mesothorax, 5-6 on metathorax, 6 on segment I. **Ducts**. Marginal macroducts, 2-barrel-shaped, 1 present between L1 and L2, 2 on segment VI, 1 on segment V. Dorsal macroducts on pygidium about same size as marginal macroducts, becoming slightly smaller on anterior abdomen, 2-barrel-shaped, arranged segmentally in submedian and submarginal rows; submedian: 3-6 on segment I, 4-5 on II, 4-6 on III, 7-13 on IV, 3-4 on V; submarginal: 11-14 on I, 11-12 on II, 10-11 on III, 10-11 on IV, 8-9 on V. Dorsal ducts scattered on margin of thorax, smaller than those on abdomen, 2-barrel-shaped, with 8 or 9 on prothorax, 15-17 on mesothorax, 14 or 15 on metathorax. Dorsal ducts on head as big as ventral microducts, very smaller than dorsal ducts present on thorax, scattered distribution. Ventral microducts scattered, numerous on head and with several microducts on submargin of pygidium and prothorax and submedian of abdomen, meso- and metathorax. **Anal opening**, small, 15-17 μm in diameter, positioned 214 μm from posterior margin. **Perivulvar pores** in 5 groups, 31-37 in the median group, 33-44 in the anterolaterally and 45-48 in the posterolaterally.

Diagnosis. This species is similar to *P. chinensis* (Cockerell, 1896) in body shape and the number of pygidial lobes, but can be distinguished by the following features (those for *P. chinensis* in brackets): 1) dorsal macroducts absent on abdominal segment VI (present); 2) L1 prominent the pygidium (sunken into the pygidium).

Host: *Spermadictyon suaveolens*.

Etymology. The specific epithet is named after Zhenyuan, the type locality.

Distribution. China (Guizhou).

Key to Chinese species of the genus *Pseudaulacaspis*

- 1 Trilocular pores absent near each anterior spiracle *P. manni* (Green & Mann, 1907)
- Trilocular pores present near each anterior spiracle 2
- 2 Body slender, both side nearly parallel *P. dendrobii* Kuwana & Muramatsu, 1931
- Body nonslender 3
- 3 Body suborbicular or oval 4
- Body long ovate or furiform 8
- 4 Trilocular pores present near each anterior spiracle, absent near each posterior spiracle 5
- Trilocular pores present near anterior spiracle and posterior spiracle 7
- 5 With 2 pairs of lobes on pygidium *P. canarium* Hu, 1986
- With 3 pairs of lobes on pygidium 6
- 6 The eggs white or salmon; with 1 pairs of gland spines between L3 and the traces of L4, each bifurcate *P. pentagona* (Targioni Tozzetti, 1886)
- The eggs always salmon; with 2 pairs of gland spines between L3 and the traces of L4, each pointed *P. prunicola* (Maskell, 1895)
- 7 Perivulvar pores in 6 groups *P. mirabilis* Hu, 1986

- Perivular pores in 5 groups *P. ficicola* Tang, 1986
- 8 Anterior spiracle and posterior spiracle both with trilocular pores 9
- Anterior spiracle with trilocular pores, posterior spiracle without trilocular pores 14
- 9 Dorsal macroducts absent on submarginal and submedial area of abdominal segment VI 10
- Dorsal macroducts present on submarginal or submedial area of abdominal segment VI 11
- 10 Submarginal and submedial macroducts present on abdominal segment I
..... *P. zhenyuanensis* sp. n.
- Submarginal and submedial macroducts absent on abdominal segment I.....
..... *P. ulmicola* Tang & Li, 1988
- 11 Dorsal macroducts absent on submarginal and submedial area of abdominal segment II..... 12
- Dorsal macroducts present on submarginal or submedial area of abdominal segment II..... 13
- 12 Antennae 6 segments in first instar *P. centreesa* (Ferris, 1953)
- Antennae 5 segments in first instar *P. eucalypticola* Tang, 1986
- 13 Dorsal macroducts present on submedial area of abdominal segment VI; anal opening situated at the base of pygidium *P. momi* (Kuwana, 1931)
- Dorsal macroducts present between submarginal or submedial area of abdominal segment VI; anal opening situated on the centre of pygidium
..... *P. loncerae* Tang, 1986
- 14 Dorsal macroducts present on submarginal or submedial area of abdominal segment VI 15
- Dorsal macroducts absent on submarginal and submedial area of abdominal segment VI 23
- 15 Dorsal macroducts present on submarginal and submedial area of abdominal segment VI 16
- Dorsal macroducts only present on submedial area of abdominal segment VI 18
- 16 L2 bilobate, each with a pair of short basal scleroses.....
..... *P. sasakawai* Takagi, 1970
- L2 bilobate, without basal scleroses..... 17
- 17 With 9-11 trilocular pores near each anterior spiracle
..... *P. camelliae* (Chen, 1983)
- With more than 25 trilocular pores near each anterior spiracle
..... *P. latisoma* (Chen, 1983)
- 18 Dorsal macroducts present on submarginal or submedial area of abdominal segment I 19
- Dorsal macroducts absent on submarginal or submedial area of abdominal segment I 20

19	L3 not obvious, anal opening situated on the centre of pygidium	<i>P. takahashii</i> (Ferris, 1955)
–	L3 bilobate, anal opening situated at the base of 2/5 of pygidium	<i>P. chinensis</i> (Cockerell, 1896)
20	L3 bilobate	21
–	L3 not obvious, present by a prominence	22
21	L1 protruding from pygidial margin	<i>P. cockerelli</i> (Cooley, 1897)
–	L1 sunk into apex of pygidium	<i>P. kentiae</i> (Kuwana, 1931)
22	Only 1 submedial macroduct present on abdominal segment VI	<i>P. eugeniae</i> (Maskell, 1892)
–	With 2 submedial macroducts present on abdominal segment VI	<i>P. ericeae</i> (Ferris, 1953)
23	Submarginal macroducts present on abdominal segment II, submedial macroducts absent on abdominal segment II	24
–	Submarginal and submedial macroducts both present on abdominal segment II	26
24	Submedial macroducts absent on abdominal segment III	<i>P. subcorticalis</i> (Green, 1905)
–	Submedial macroducts present on abdominal segment III	25
25	With more than 30 trilocular pores near each anterior spiracle	<i>P. poloosta</i> (Ferris, 1953)
–	With 20 or fewer trilocular pores near each anterior spiracle	<i>P. megaloba</i> (Green, 1899)
26	L3 obvious, bilobate	27
–	L3 not obvious, present by a shallow prominence	28
27	With 2-4 trilocular pores near each anterior spiracle	<i>P. subrhombica</i> (Chen, 1983)
–	With 11-14 trilocular pores near each anterior spiracle	<i>P. frutescens</i> (Hu, 1986)
28	With 10 or fewer trilocular pores near each anterior spiracle	29
–	With more than 15 trilocular pores near each anterior spiracle	31
29	L1 protruding from pygidial margin, only with 2 trilocular pores near each anterior spiracle	<i>P. taiwana</i> (Takahashi, 1935)
–	L1 sunk into apex of pygidium, with more than 4 trilocular pores near each anterior spiracle	30
30	The terminal of L3 arc-shaped, smoothly, with 4-8 trilocular pores near each anterior spiracle	<i>P. abbrideliae</i> (Chen, 1983)
–	The terminal of L3 serration, with more than 4 trilocular pores near each anterior spiracle	<i>P. brideliae</i> (Takahashi, 1933)
31	Gland tubercles present on prothorax, dorsal ducts present on head, smaller than those present on pygidium and abdomen....	<i>P. syzygicola</i> (Tang, 1986)
–	Gland tubercles absent on prothorax, dorsal ducts absent on head	32

- 32 L2 small, bilobulate, with 17–20 submedial macroduct in total.....
 *P. kuishiensis* (Kuwana, 1909)
- L2 very small, the outer lobule at times almost obsolete, with 4–11 submedial
 macroduct in total *P. celtis* (Kuwana, 1928)

Acknowledgements

This study is supported by the National Natural Science Foundation of China (Grant No. 30870324).

Reference

- Andersen JC, Wu J, Gruwell ME, Gwiazdowski R, Santana SE, Feliciano NM, Morse GE, Normark BB (2010) A phylogenetic analysis of armored scale insects (Hemiptera: Diaspididae), based upon nuclear, mitochondrial, and endosymbiont gene sequences. *Molecular Phylogenetics and Evolution* 57: 992–1003. doi: 10.1016/j.ympev.2010.05.002
- Ben-Dov Y (2012) Scalenet. Species in genus *Pseudaulacaspis* query result. <http://scalenet.info/validname/Pseudaulacaspis/> [Accessed February 2012]
- Borchsenius NS (1966) A catalogue of the armoured scale insects (Diaspidoidea) of the world. Nauka, Moscow, Leningrad, 449 pp. [In Russian]
- Chen FG (1983) The Chionaspini (Diaspididae, Coccoidea, Homoptera) from China. Science & Technology Publishing House, Sichuan Province, 175 pp.
- Ferris GF (1953) Report upon scale insects collected in China (Homoptera: Coccoidea). Part IV. (Contribution No. 84). *Microentomology* 18: 59–84.
- Ferris GF (1955) The genus *Phenacaspis* Cooley and Cockerell, Part I. (Insecta: Homoptera: Coccoidea) (Contribution No. 93). *Microentomology* 20: 41–82.
- Gullan PJ, Cook LG (2007) Phylogeny and higher classification of the scale insects (Hemiptera: Sternorrhyncha: Coccoidea). *Zootaxa* 1668: 413–425.
- Henderson RC (2011) Diaspididae (Insecta: Hemiptera: Coccoidea). *Fauna of New Zealand* 66. Manaaki Whenua Press, Lincoln, Canterbury, 275 pp.
- Hodgson CJ, Lagowska B (2011) New scale insect (Hemiptera: Sternorrhyncha: Coccoidea) records from Fiji: three new species, records of several new invasive species and an updated checklist of Coccoidea. *Zootaxa* 2766: 1–29.
- Hu JL (1986) Studies on scale insects in the Hainan Island of China (Part I). *Contributions from Shanghai Institute of Entomology* 6: 213–227.
- MacGillivray AD (1921) The Coccidae. Tables for the Identification of the Subfamilies and Some of the More Important Genera and Species, Together with Discussions of Their Anatomy and Life History. Scarab, Urbana, Ill., 502 pp.
- Miller DR, Davidson JA (2005) Armored Scale Insect Pests of Trees and Shrubs. Cornell University Press, Ithaca, 442 pp.

- Morse GE, Normark BB (2006) A molecular phylogenetic study of armoured scale insects (Hemiptera: Diaspididae). *Systematic Entomology* 31: 338–349. doi: 10.1111/j.1365-3113.2005.00316.x
- Takagi S (1956) Notes on the Japanese species of the genus *Pseudaulacaspis* MacGillivray, with description of a new species. *Insecta Matsumurana* 19: 113–116.
- Takagi S (1961) A contribution to the knowledge of the Diaspididini of Japan (Homoptera: Coccoidea) Pt. III. *Insecta Matsumurana* 24: 69–103.
- Takagi S, Kawai S (1966) Some Diaspididae of Japan (Homoptera: Coccoidea). *Insecta Matsumurana* 28: 93–119.
- Takagi S (1970) Diaspididae of Taiwan based on material collected in connection with the Japan-US Cooperative Science Programme, 1965 (Homoptera: Coccoidea). Pt. II. *Insecta Matsumurana* 33: 1–146.
- Takagi S (1985) The scale insect genus *Chionaspis*: A revised concept (Homoptera: Coccoidea: Diaspididae). *Insecta Matsumurana Series Entomology. New Series*, 33: 1–77.
- Tang FT (1986) The Scale Insects of Horticulture and Forests of China. Volume III. Shanxi Agricultural University Press, Taigu, Shanxi, 305 pp.
- Tang FT, Li J (1988) Observations on the Coccoidea of Inner Mongolia in China. Inner Mongolia University Press, 227 pp.
- Williams DJ, Watson GW (1988) The Scale Insects of the Tropical South Pacific Region. Pt. 1. The Armoured Scales (Diaspididae). CAB International Institute of Entomology, London, 290 pp.