

# A new species of *Parlatoria* from China (Hemiptera, Coccoomorpha, Diaspididae)

Minmin Niu<sup>1</sup>, Jinian Feng<sup>1</sup>

<sup>1</sup> Key Laboratory of Plant Protection Resources and Pest Management, Ministry of Education, Entomological Museum, College of Plant Protection, Northwest A&F University, Yangling, Shaanxi Province, 712100, China

Corresponding author: Jinian Feng ([jinianf@nwfau.edu.cn](mailto:jinianf@nwfau.edu.cn))

---

Academic editor: R. Blackman | Received 14 June 2018 | Accepted 8 July 2018 | Published 7 August 2018

<http://zoobank.org/C2DD2540-4FB5-433E-A625-41261CC3E4D8>

---

**Citation:** Niu M, Feng J (2018) A new species of *Parlatoria* from China (Hemiptera, Coccoomorpha, Diaspididae). ZooKeys 779: 147–155. <https://doi.org/10.3897/zookeys.779.27437>

---

## Abstract

A new species of armoured scale insect, *Parlatoria menglaensis* sp. n. is described and illustrated, which infests leaves of *Cinnamomum camphora* in China. A key to the *Parlatoria* species occurring in China is provided.

## Keywords

China, Diaspididae, Hemiptera, new species, *Parlatoria*

## Introduction

Armoured scale insects (Hemiptera: Coccoomorpha: Diaspididae) are ubiquitous sap-sucking parasites that have a worldwide distribution (Andersen 2009). Compared with other family groups of Coccoidea, the armoured scale insects have several unique characteristics. For example, they have a cecum, which has no direct connection between the stomach and the anal opening, so they do not produce sweet secretions called honeydew (Henderson 2011). Females of the armoured scale insects are immobile and firmly attached to plant leaves, stems, fruits, or roots (Chou 1982). They have an extremely simplified morphology with several fused segments (Andersen 2010). They have no wings, or legs, and the eyes and antennae are reduced (Balachowsky 1948). The Diaspididae is the largest family of the Coccoidea with 2595 species currently identified (García Morales et al. 2018).

The genus, *Parlatoria* Targioni Tozzetti, 1868, is a large group of Diaspididae that are members of the tribe Parlatoriini. This genus was originally established by Curtis (1843) and *Aspidiotus proteus* Curtis, 1843, has been subsequently designated as the type species by Leonardi. The genus currently is made up of 73 species (García Morales et al. 2018). Some species of *Parlatoria*, such as *P. oleae* (Colvée) and *P. ziziphi* (Lucas) are considered to be serious pests of economic plants (McKenzie 1945).

*Parlatoria* is distributed in both tropical and subtropical regions (García Morales et al. 2018). This genus is found predominately in southeastern Asia but has also extended its range into Australia and Africa (McKenzie 1945). Some species have been introduced in the tropics and other warm parts of the world (Takagi 1969). About 34 species of this genus have been reported from China (García Morales et al. 2018).

Recently, a new species of *Parlatoria* was discovered in China and is described and illustrated in this study. This discovery raises the number of species recorded in this genus to 74, of which 35 are recorded from China. A key to the Chinese species of *Parlatoria* is presented in this study.

## Materials and methods

Plant samples infested by the new species were collected from Mengla city, Yunnan Province. Permanent slide mounts of adult females from the samples were prepared using the method described by Henderson (2011).

The illustrations of the adult female shown in Figs. 1–9 were drawn from slide-mounted specimens. Fig. 1 shows an overview of the dorsal body surface on the left side and the ventral body surface on the right side, and an enlarged detail of the significant features of the body, which are not drawn in direct proportions to each other.

Slide-mounted type specimens of the new species have been deposited in the Entomological Museum, Northwest A&F University, Yangling, Shaanxi, China (NWFU).

## Taxonomy

### *Parlatoria* Targioni Tozzetti

*Parlatoria* Targioni Tozzetti, 1868: 735.

**Type species.** *Parlatoria orbicularis* Targioni Tozzetti, subsequently designated by Leonardi (1899: 208).

**Generic diagnosis. Female scale.** Various colors, oval to elongate, with exuviae terminal, often occupying greater part of scale.

**Male scale.** Elongate and smaller than the female.

**Adult female.** Female body nearly round or broadly ovate, and the widest at meta-thorax or abdominal segment I. The body free segments distinct but not strongly pro-

duced, and membranous except for the pygidium. Antennae with one or two setae. Prosomatic tubercles present or absent. Anterior spiracles with disc pores, the posterior spiracles without disc pores. Derm pocket present or absent between the posterior spiracle and body margin. Peribuccal granulations present or absent. The pygidium rounded or rather triangular. Three pairs of lobes well developed, unilobate, usually notched, median lobes not yoked; marginal macroducts of pygidium, one present or absent between the median lobes, one between the median and second lobes, and also between the second and third lobes. Two or three fimbriate plates present between lobes, then extend to the whole lateral margin of the pygidium, and each with one microduct. The fourth and fifth lobes often present, sclerotized or replaced by a membranous fimbriate plates. Submarginal dorsal ducts usually present, scattered in a broad, continuous, irregular row on each side of the abdomen. Submedian dorsal ducts present or absent. Anal opening positioned about centre of pygidium. Four or five groups of perivulvar pores present (Adapted from McKenzie 1945, Takagi 1969, Williams and Watson 1988, Henderson 2011).

**Remarks.** This genus *Parlatoria*, like other groups of the subfamily Aspidiotinae has an ovate body, and the second lobes are not divided into two lobules. Fringed plates are present between the lobes. *Parlatoria* is distinguished from other genera, especially *Parlagena* McKenzie, 1945, and *Parlatoreopsis* Lindinger, 1912, by having fringed plates across two sides of the prepygidial abdominal segments.

***Parlatoria menglaensis* sp. n.**

<http://zoobank.org/5F9869BC-25A3-4633-B445-6C6B76453DD8>

Figures 1–9


**Material examined.** Holotype and 30 paratypes, adult female. China: Yunnan Province. Mengla city, longitude 101.57, latitude: 21.48, on *Cinnamomum camphora* (L.) Presl., 30.vii.2017, coll. Minmin Niu.

**Description.** Female scale. Adult female cover convex, circular dark green/grey; exuvia on front end. Male scale. Not recorded.

**Adult female.** Body elongate-ovate, the broadest in first abdominal segment or thoracic region; segmentation distinct.

Antenna composed of two stout tubercles and a seta, located midway between frontal margin and mouthparts, interantennal distance being equal to width of mouthparts. Prosomatic tubercles absent. Anterior spiracle with 1–2 parastigmatic pores; posterior spiracle without pores. Derm pocket absent between the posterior spiracle and body margin. Submarginal duct tubercles arranged as follows: 1–2 on cephaloprothoracic sternum, 3–5 on the mesothorax, and 4–8 on the metathorax.

Pygidium rounded, with 3 pairs of well-developed lobes. The median lobes distinctly notched once on each side, the second and third usually notched once distinctly on outer side. Fourth lobes replaced by a fimbriate plate, similar to adjacent fimbriate plates in form. Fimbriate plates present throughout pygidium, mostly as long as lobes, and two between median lobes, two between the median and second, three between



**Figure 1–9.** *Parlatoria menglaensis* sp. n., adult female. **1** body **2** antenna **3** anterior spiracle **4** posterior spiracle **5** gland tubercle **6** detail of dorsal gland duct **7** ventral microduct **8** perivulvar pore **9** pygidium.

the second and third; pygidial marginal macroducts largest closer to median lobes, arranged as follows: absent between the median lobes, one in each interlobar space, two outside the third lobe. Submarginal dorsal ducts on prepygidium numerous, 60–70 ducts on each side. Submedian ducts absent. Ventral microducts few, scattered on pygidium. Anal opening small, positioned about centre of pygidium. Perivulvar pores present in five groups; 3–4 in the median group, laterocephalic group with four pores, and the laterocaudal group with four pores.

**Remarks.** This species is very similar to *P. machilicola* (Takahashi 1933) in body shape, but differs in having (character-states on *P. machilicola* in brackets): (i) perivulvar pores present in five groups (perivulvar pores present in four groups); (ii) dorsal macroducts absent (dorsal macroducts present); (iii) marginal macroducts only four on each side (marginal macroducts more than four on each side).

The new species also resembles *P. tsujii* Tanaka, 2010, in the shape and spiracles. However, it differs from *P. tsujii* by the following characters (character-states on *P. machilicola* in brackets): (i) perivulvar pores present in five groups (perivulvar pores present in four groups); (ii) derm pocket absent (derm pocket present); (iii) marginal macroducts absent between the median lobes (marginal macroducts present between the median lobes).

**Host plant.** *Cinnamomum camphora* (L.) Presl (Lauraceae).

**Etymology.** The specific epithet is named after Mengla, the type locality.

**Distribution.** China (Yunnan).

### Key to the adult females of *Parlatoria* Targioni Tozzetti from China

(There are records of *P. ligustri* Wu from China (Wu Chenfu 1935), but the information is inadequate for this key, thus it has not been included).

- |   |                                                                                                                                                                                        |                                       |
|---|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|
| 1 | Perivulvar pores absent.....                                                                                                                                                           | <i>P. pseudaspidiotus</i> (Lindinger) |
| – | Perivulvar pores present in 4 or 5 groups.....                                                                                                                                         | 2                                     |
| 2 | A marginal macroduct absent between the median lobes .....                                                                                                                             | 3                                     |
| – | A marginal macroduct present between the median lobes .....                                                                                                                            | 4                                     |
| 3 | Perivulvar pores absent in the median group .....                                                                                                                                      | <i>P. machilicola</i> (Takahashi)     |
| – | Perivulvar pores present in the median group.....                                                                                                                                      | <i>P. menglaensis</i> sp. n.          |
| 4 | Puparium of adult female black; with 1 peculiar, large and ear-like lobe on each side margin of head region about opposite anterior spiracles .....                                    | <i>P. ziziphi</i> (Lucas)             |
| – | Puparium of adult female variable in color, but not black; on margin without ear-like lobes or lobes very small on each side of head region about opposite of anterior spiracles ..... | 5                                     |
| 5 | Pygidium with 2 pairs of well-developed lobes, the 3 <sup>rd</sup> quite small, but sclerotized .....                                                                                  | <i>P. cupressi</i> (Ferris)           |
| – | Pygidium with 3 or more pairs of well-developed lobes.....                                                                                                                             | 6                                     |

6	Anterior spiracles without disc pores.....	<i>P. mytilaspiformis</i> Green
–	Anterior spiracles with disc pores.....	7
7	Derm pockets present between each posterior spiracle and body margin.....	8
–	Derm pockets absent between each posterior spiracle and body margin....	13
8	Fourth lobes definitely present, not closely resembling adjacent plates.....	9
–	Fourth lobes absent, replaced by a membranous, plate-like process, smaller than the adjacent plates.....	<i>P. proteus</i> (Curtis)
9	Four plates present between third lobe and fourth lobe.....	<i>P. pinicola</i> Tang
–	Three plates present between third lobe and fourth lobe.....	10
10	Eyespots modified to form a stout spur.....	<i>P. crotonis</i> (Douglas)
–	Eyespots various, flat, irregular or absent.....	11
11	Peribuccal granulations absent.....	<i>P. camelliae</i> (Comstock)
–	Peribuccal granulations present.....	12
12	Pygidial lobes almost equal in size; perivulvar pores always in 4 groups.....	<i>P. pini</i> Tang
–	Pygidial lobes of different sizes, the third pair smallest; perivulvar pores sometimes present in 5 groups.....	<i>P. theae</i> (Cockerell)
13	Pygidium normally with four plates present between third lobe and fourth lobe.....	14
–	Pygidium with three plates present between third lobe and fourth lobe....	17
14	Peribuccal granulations present.....	15
–	Peribuccal granulations absent.....	<i>P. multipora</i> McKenzie
15	Dorsal ducts present on submedian area on abdominal segments I-III.....	<i>P. yanyuanensis</i> Tang
–	Dorsal ducts absent from submedian area on abdominal segments I-III....	16
16	Median lobes usually with a deep lateral notch and no medial notch (rarely with a small medial notch).....	<i>P. oleae</i> (Colvée)
–	Median lobes with deep lateral and medial notches.....	<i>P. bambusae</i> Tang
17	Fourth lobes definitely present, not closely resembling adjacent plates.....	18
–	Fourth lobes replaced by a membranous, plate-like process, smaller than the adjacent plates.....	27
18	Lobes with outer margins normally notched.....	19
–	Lobes with outer margins minutely toothed.....	<i>P. yunnanensis</i> Ferris
19	Lobes with outer margins with more than one notch.....	20
–	Lobes with outer margins with a single notch.....	22
20	Plates reduced.....	21
–	Plates well developed, fimbriate.....	<i>P. desolator</i> McKenzie
21	Dorsal median macroducts present on pygidium within frame formed by perivulvar pores.....	<i>P. cinerea</i> (Doane & Hadden)
–	Dorsal median macroducts absent on pygidium within frame formed by perivulvar pores.....	<i>P. fluggeae</i> Hall
22	Peribuccal granulations present.....	23
–	Peribuccal granulations absent.....	24

- 23 Submedian dorsal macroducts absent; Eyespot modified to form rounded elevation..... *P. machili* Takahashi
- Submedian dorsal macroducts present on abdominal segments IV–V; Eyespot absent or inconspicuous ..... *P. cinnamomicola* Tang
- 24 Median pygidial lobes deeply notched on outer and inner margins ..... 25
- Median pygidial lobes deeply notched only on outer margins ..... 26
- 25 Perivulvar pores absent in the median group, antero- and posterolateral groups confluent on each side; submedian dorsal macroducts present on abdominal segments I–III ..... *P. emeiensis* Tang
- Perivulvar pores absent or a single one present in the median group, 6 to 8 in the anterolateral group, and 6 to 8 in the posterolateral group; submedian dorsal macroducts absent ..... *P. pergandii* (Comstock)
- 26 Fourth lobes represented by small sclerotized spurs ...*P. ghanii* Hall & Williams
- Fourth lobes well developed..... *P. reedia* Zhang, Feng & Liu
- 27 Peribuccal granulations absent ..... 28
- Peribuccal granulations present ..... 30
- 28 Anterior spiracle with 1–3 disc pores ..... 29
- Anterior spiracle with 6–15 disc pores ..... *P. stigmadisculosa* Bellio
- 29 Plates tapering apically to a point..... *P. lithocarpi* Takahashi
- Plates well fimbriated..... *P. acalcarata* McKenzie
- 30 Perivulvar pores present in the median group, with 5 groups ...*P. piniphila* Tang
- Perivulvar pores absent in the median group ..... 31
- 31 Prosomatic tubercles rounded ..... *P. arengae* Takagi
- Prosomatic tubercles pointed apically, inconspicuous or absent..... 32
- 32 Posterolateral groups of perivulvar pores with 2–3 pores each ..... *P. hydnocarpus* Hu
- Posterolateral groups of perivulvar pores with more than 6 pores each ..... 33
- 33 Gland tubercles: 1 to 3 prespiraculars, 3 to 4 anterior spiraculars, 3 to 5 mesothoracics, 4 metathoracics, and 4 first abdominals; submarginal dorsal ducts containing 18 to 22 on each side ..... *P. liriopicola* Tang
- Gland tubercles: 1 to 7 prespiraculars, 4 to 10 anterior spiraculars, 14 to 16 mesothoracics, 8 to 16 metathoracics, and 5 to 8 first abdominals; submarginal dorsal ducts containing 35 to 47 on each side ..... *P. keteleericola* Tang & Chu

## Discussion

*Parlatoria menglaensis* sp. n. is a pest in urban areas of China. Its primary host plant is *Cinnamomum camphora* (L.) Presl (Lauraceae), which grows south of the Yangtze River in China. This plant is also found in southern Japan, Korea, and Vietnam, and has been introduced to many other countries, including Australia and the United States. Currently, *P. menglaensis* sp. n. has only been identified in China, but it could eventually spread to the previously mentioned countries where its potential host plants occur.



## Acknowledgements

We are grateful to Prof. William H. Reissig from Cornell University (New York, USA) for language revision of the manuscript. The work is supported by the National Natural Science Foundation of China (No. 31472022 and No. 31772502).

## References

- Andersen JC (2009) A phylogenetic analysis of armored scale insects, based upon nuclear, mitochondrial, and endosymbiont gene sequences. University of Massachusetts Amherst, 66 pp.
- Andersen JC, Wu J, Gruwell ME, Morse GE, Santana S, Feliciano N, Gwiazdowski RA, Normark BB (2010) A phylogenetic analysis of armored scale insects, based up on nuclear, mitochondrial, and endosymbiont gene sequences. *Molecular Phylogenetics and Evolution* 57: 992–1003. <https://doi.org/10.1016/j.ympev.2010.05.002>
- Balachowsky AS (1948) Les cochenilles de France, d'Europe, du nord de l'Afrique et du bassin Méditerranéen. IV. Monographie des Coccoïdea, classification – Diaspidinae (Première partie). *Actualités Scientifiques et Industrielles* 1054: 243–394.
- Bellio G (1929) Note su Diaspinae dell' estremo oriente con descrizione di nuove specie (Homoptera: Coccidae). *Bollettino del R. Istituto Superior Agraria Laboratorio di Zoologia Generale e Agraria* 22: 219–243.
- Chou I (1982) Monograph of the Diaspididae of China. Vol. 1. Shanxi Publication House of Science & Technology Shanxi, 195 pp.
- Cockerell TDA (1896) Some coccidae found by Mr. Alex Craw in the course of his quarantine work at San Francisco. United States Department of Agriculture, Division of Entomology, Technical Series 4: 42–46.
- Colvée P (1880) Ensayo sobre una nueva enfermedad del olivo, producida por una nueva especie del género *Aspidiotus*. *Gaceta Agrícola del Ministerio de Fomento* 14: 21–41.
- Comstock JH (1881) Report of the Entomologist. Report of the Commissioner of Agriculture, United States Department of Agriculture 1880/1881: 276–349.
- Comstock JH (1883) Second report on scale insects, including a monograph of the sub-family Diaspinae of the family Coccidae and a list, with notes of the other species of scale insects found in North America. Department of Entomology Report, Cornell University Agricultural Experiment Station 2: 47–142.
- Curtis J (1843) The small brown scale, *Aspidiotus proteus*, Nobis. *Gardeners' Chronicle* 39: 676.
- Doane RW, Hadden E (1909) Coccidae from the Society Islands. *Canadian Entomologist* 41: 296–300. <https://doi.org/10.4039/Ent41296-8>
- Douglas JW (1887) Note on some British Coccidae (No. 6). *Entomologist's Monthly Magazine* 23: 239–243. <https://doi.org/10.5962/bhl.part.18592>
- Ferris GF (1953) Report upon scale insects collected in China (Homoptera: Coccoidea). Part IV. (Contribution No. 84). *Microentomology* 18: 59–84.
- García Morales M, Denno B, Miller DR, Miller GL, Ben-Dov Y, Hardy NB (2018) ScaleNet: A Literature based model of scale insect biology and systematics. <http://scalenet.info>



- Green EE (1899) The Coccidae of Ceylon, Part II. Dulau London, 105–169.
- Hall WJ (1929) Observations on the Coccidae of southern Rhodesia. III. Bulletin of Entomological Research 20: 359–376. <https://doi.org/10.1017/S0007485300021313>
- Hall WJ, Williams DJ (1962) New Diaspididae (Homoptera: Coccoidea) from the Indo-Malayan region. Bulletin of the British Museum (Natural History) Entomology 13: 21–43.
- Henderson RC (2011) Diaspididae (Insecta: Hemiptera: Coccoidea). Fauna of New Zealand 66. Manaaki Whenua Press, Lincoln, Canterbury, 275 pp.
- Hu J (1986) Studies on scale insects in the Hainan Island of China (Part I). Contributions from Shanghai Institute of Entomology 6: 213–227.
- Leonardi G (1899) Sistema delle “Parlatoriae” nota preventiva. Mariani Florence, 203–209.
- Lindinger L (1905) Zwei neue Schildläuse aus Asien. Insecten Börse 22: 131–132.
- Lindinger L (1912) Die Schildläuse (Coccidae) Europas, Nordafrikas und Vorder-Asiens, einschliesslich der Azoren, der Kanaren und Madeiras. Ulmer, Stuttgart, 388 pp.
- Lucas H (1853) [*Coccus ziziphi*]. Annales de la Société Entomologique de France 1: 28–29.
- McKenzie HL (1945) A revision of *Parlatoria* and closely allied genera (Homoptera: Coccoidea: Diaspididae). Microentomology 10: 47–121.
- McKenzie HL (1960) Taxonomic position of *Parlatoria virescens* Maskell, and descriptions of related species (Homoptera: Coccoidea: Diaspididae). Scale studies pt. 14. Bulletin of the California Department of Agriculture 49: 204–211.
- Takagi S (1969) Diaspididae of Taiwan based on material collected in connection with the Japan-U.S. Co-operative Science Programme, 1965 (Homoptera: Coccoidea). Part I. Insecta Matsumurana 32: 1–110.
- Takahashi R (1931) Descriptions of some new Formosan Coccidae. (Rhynchota). Bulletin of Entomological Research 22: 211–220. <https://doi.org/10.1017/S0007485300035173>
- Takahashi R (1933) Observations on the Coccidae of Formosa. III. Report. Government Research Institute. Department of Agriculture. Formosa 60: 1–64.
- Takahashi R (1935) Observations on the Coccidae of Formosa. V. Report Department of Agriculture Government Research Institute, Formosa 66: 1–37.
- Tanaka H (2010) A New Species of the Genus *Parlatoria* (Hemiptera, Diaspididae) from Japan. Japanese Journal of Systematic Entomology 16(2): 179–183.
- Tang FT (1984) The scale insects of horticulture and forests of China. Shanxi Agricultural University Press Research Publication 2: 1–115.
- Tang FT, Chu CY (1983) Three new species of Diaspididae from *Keteleeria evelyniana* in vicinity of Kunming, Yunnan. Acta Zootaxonomica Sinica 8: 301–306.
- Targioni Tozzetti A (1868) Introduzione alla seconda memoria per gli studi sulle cocciniglie, e catalogo dei generi e delle specie della famiglia dei coccidi. Atti della Società italiana di scienze naturali 11: 721–738.
- Williams DJ, Watson GW (1988) The Scale Insects of the Tropical South Pacific Region. Pt. 1: The Armoured Scales (Diaspididae). CAB International Wallingford, 290 pp.
- Wu Chenfu F (1935) Catalogus Insectorum Sinensium (catalogue of Chinese insects). Volume II. The Fan Memorial Institute of Biology, Peiping, China, 639 pp.
- Zhang FP, Feng JN, Liu XW (2006) A new species and a new record of *Parlatoria* from China (Hemiptera: Diaspididae). Acta Zootaxonomica Sinica 31(4): 832–834.